

7. Sabour H, Javidan AN, Vafa MR, et al. Calorie and macronutrients intake in people with spinal cord injuries: an analysis by sex and injury-related variables. *Nutrition* 2012;28:143-7.
8. Turner JE, Markovitch D, Betts JA, Thompson D. Nonprescribed physical activity energy expenditure is maintained with structured exercise and implicates a compensatory increase in energy intake. *Am J Clin Nutr* 2010;92:1009-16.
9. Fontana L, Villareal DT, Weiss EP, et al. Calorie restriction or exercise: effects on coronary heart disease risk factors. A randomized, controlled trial. *Am J Physiol Endocrinol Metab* 2007;293:E197-202.
10. Naghavi M, Libby P, Falk E, et al. From vulnerable plaque to vulnerable patient: a call for new definitions and risk assessment strategies: Part I. *Circulation* 2003;108:1664-72.
11. Mora S, Cook N, Buring JE, Ridker PM, Lee IM. Physical activity and reduced risk of cardiovascular events: potential mediating mechanisms. *Circulation* 2007;116:2110-8.
12. Green DJ, O'Driscoll G, Joyner MJ, Cable NT. Exercise and cardiovascular risk reduction: time to update the rationale for exercise? *J Appl Physiol* 2008;105:766-8.
13. Green DJ, Maiorana A, O'Driscoll G, Taylor RR. Topical review: effects of exercise training on vascular endothelial nitric oxide function in humans. *J Physiol* 2004;561:1-25.
14. Dinunno FA, Tanaka H, Monahan KD, et al. Regular endurance exercise induces expansive arterial remodelling in the trained limbs of healthy men. *J Physiol* 2001;534:287-95.
15. Levy WC, Cerqueira MD, Harp GD, et al. Effect of endurance exercise training on heart rate variability at rest in healthy young and older men. *Am J Cardiol* 1998;82:1236-41.
16. Tsuji H, Venditti Jr FJ, Manders ES, et al. Reduced heart rate variability and mortality risk in an elderly cohort. The Framingham Heart Study. *Circulation* 1994;90:878-83.
17. Latimer AE, Ginis KA, Craven BC, Hicks AL. The physical activity recall assessment for people with spinal cord injury: validity. *Med Sci Sports Exerc* 2006;38:208-16.
18. Dwyer TJ, Alison JA, McKeough ZJ, Elkins MR, Bye PT. Evaluation of the SenseWear activity monitor during exercise in cystic fibrosis and in health. *Respir Med* 2009;103:1511-7.

<http://dx.doi.org/10.1016/j.apmr.2016.02.021>

Corrections



In the article "Exercise Recommendations and Considerations for Persons With Spinal Cord Injury" (Evans et al, *Arch Phys Med Rehabil* 2015;96:1749-50), these cardiovascular and muscular strength/endurance recommendations are a modified version of the Physical Activity Guidelines for Adults with Spinal Cord Injury developed and published by Martin Ginis KA, Hicks AL, Latimer AE, et al. The development of evidence-informed physical activity guidelines for adults with spinal cord injury. *Spinal Cord* 2011;49:1088-96.

0003-9993/16

<http://dx.doi.org/10.1016/j.apmr.2016.03.005>



In the article "Timed 360° Turn Test for Assessing People With Chronic Stroke" (Shiu et al, *Arch Phys Med Rehabil* 2016;97:536-44), the sentence in the abstract that reads Participants: Individuals with chronic stroke (n=72) and healthy individuals (n=35) of similar age (N=107).

Should read:

Participants: Individuals with chronic stroke (n=37) and healthy individuals (n=35) of similar age (N=72).

0003-9993/16

<http://dx.doi.org/10.1016/j.apmr.2016.04.003>
