

DEPARTMENTS

Correction



In the article “Comparison of removable rigid dressing and elastic bandage for residual limb maturation in transtibial amputees: a randomized controlled trial,” published in the October 2020 issue (VOLUME 101, ISSUE 10, P1683-1688, DOI: <https://doi.org/10.1016/j.apmr.2020.05.009>, [https://www.archives-pmr.org/article/S0003-9993\(20\)30298-7/fulltext](https://www.archives-pmr.org/article/S0003-9993(20)30298-7/fulltext), there were errors in Table 1. Some data were switched between two columns. The corrected table is shown here.

Table 1 Characteristics of the participants

Characteristics	RRD Group(n=13)	EB Group(n=12)	P-Value*
Age (years), mean (SD)	62.15 (17.32)	65.25 (14.21)	0.632
Sex: n (%)			0.688
male	9 (69%)	7 (58%)	
female	4 (31%)	5 (42%)	
Cause of amputation: n (%)			1.000
Infected diabetes mellitus foot	7 (54%)	6 (50%)	
Critical limb ischemia	5 (38%)	6 (50%)	
Trauma	1 (8%)	0 (0%)	
Duration after amputation (days), mean (SD)	34.62 (16.51)	24.83 (13.60)	0.121
Baseline residual limb volume (cm ³), mean (SD)	894.78 (660.85)	830.92 (305.21)	0.763

* Fisher’s exact tests for categorical and Student’s t-tests for continuous variables