

## ORIGINAL RESEARCH

## Measuring Mechanical Properties of Spastic Muscles After Stroke. Does Muscle Position During Assessment Really Matter?

María-Isabel García-Bernal, MHS,<sup>a</sup> Paula González-García, PhD,<sup>a</sup>  
María Jesús Casuso-Holgado, PhD,<sup>a,b</sup> María Dolores Cortés-Vega, PhD,<sup>a</sup>  
Alberto Marcos Heredia-Rizo, PhD<sup>a,b</sup>

From the <sup>a</sup>Departamento de Fisioterapia, Facultad de Enfermería, Fisioterapia y Podología, Universidad de Sevilla, Sevilla, Spain; and <sup>b</sup>UMSS Research Group, Universidad de Sevilla, Sevilla, Spain.

**Abstract**

**Objective:** To investigate the influence of muscle position (relaxed vs stretched) on muscle mechanical properties and the ability of myotonometry to detect differences between sides, groups, and sites of testing in patients with stroke. We also analyzed the association between myotonometry and clinical measures of spasticity.

**Design:** Cross-sectional study.

**Setting:** Outpatient rehabilitation units including private and public centers.

**Participants:** Seventy-one participants (20 subacute stroke, 20 chronic stroke, 31 controls) were recruited (N=71).

**Intervention:** Muscle mechanical properties were measured bilaterally with a MyotonPRO at muscle belly and musculotendinous sites during 2 protocols (muscle relaxed or in maximal bearable stretched position).

**Main Outcome Measures:** Muscle tone and stiffness of the biceps brachii and gastrocnemius. Poststroke spasticity was evaluated with the Modified Tardieu Scale (MTS). A mixed-model analysis of variance was used to detect differences in the outcome measures.

**Results:** The analysis of variance showed a significant effect of muscle position on muscle mechanical properties (higher tone and stiffness with the muscle assessed in stretched position). Measurements with the stretched muscle could help discriminate between spastic and nonspastic sides, but only at the biceps brachii. Overall, there was a significant increase in tone and stiffness in the chronic stroke group and in myotendinous sites compared with muscle belly sites (all,  $P < .05$ ). No correlations were found between myotonometry and the MTS.

**Conclusions:** Myotonometry assessment of mechanical properties with the muscle stretched improves the ability of myotonometry to discriminate between sides in patients after stroke and between people with and without stroke.

Archives of Physical Medicine and Rehabilitation 2022;000:1–7

© 2022 by the American Congress of Rehabilitation Medicine. Published by Elsevier Inc. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>)

Spasticity is a frequent and disabling poststroke sequela, with an estimated prevalence of 25%.<sup>1</sup> Despite being a well-known disorder, there is little consensus on how to measure spasticity.<sup>2</sup> Subjective scales are common in the clinical setting, with limited evidence to support their use because they lack proper validity,<sup>3</sup> reliability, and reproducibility.<sup>4</sup> Clinical measures cannot

discriminate between the neural and nonneural (peripheral) components of spasticity,<sup>4</sup> except the Modified Tardieu Scale (MTS). The peripheral contribution to poststroke spasticity (PSS) can be quantified for clinical and research purposes using objective, non-invasive methods, for example, shear-wave elastography and myotonometry.<sup>4,5</sup> Myotonometry represents a valid, reliable, and convenient tool<sup>4,6</sup> that has proven to be useful to monitor PSS after conservative or invasive treatments.<sup>7</sup> However, current evidence on the ability of myotonometry to discriminate between spastic and nonspastic muscles after stroke is scarce and conflicting.<sup>6</sup> It

This research was partially funded by the Ilustre Colegio Profesional de Fisioterapeutas de Andalucía (grant reference no. 03729/19D/MA)  
Clinical Trial Registration No.: NCT03814460.  
Disclosures: none.

has been recommended to conduct myotonometry measurement of PSS at several muscle sites of testing and in different muscle positions, that is, relaxed or stretched,<sup>6</sup> to get a clear picture of how muscle mechanical properties may change after stroke<sup>8</sup> and in response to rehabilitation programs.

This study aimed to investigate the differences in myotonometry scores for muscle tone and stiffness in patients with stroke, comparing sides (affected vs nonaffected), sites (muscle belly [MB] vs musculotendinous [MT]), and groups (subacute stroke, chronic stroke, control), during 2 evaluation protocols (relaxed or stretched muscle). As a secondary goal, we analyzed the possible associations between myotonometry and the MTS. We hypothesized differences between protocols in myotonometry scores and that measuring tone and stiffness in stretched position would help to better distinguish between the affected and nonaffected sides in patients with stroke and between individuals with or without stroke.

## Methods

### Design

We conducted a multicenter, cross-sectional study, including adults with subacute (6-36 weeks after the event)<sup>9</sup> or chronic (more than 36 weeks) stroke<sup>10</sup> and participants without stroke. The protocol of the study respected the ethical guidelines set in the Helsinki Declaration and was approved by the Junta de Andalucía Ethical Committee for Biomedical Research (CI 1222-N-16). It followed the Strengthening the Reporting of Observational Studies in Epidemiology framework for observational studies. All participants provided verbal and written informed consent.

### Participants

Individuals with a first-ever stroke were selected from public and private centers. Participants should have at least a slight increase of biceps brachii and gastrocnemius muscle tone. This was identified with a score  $\geq 1$  in the Modified Ashworth Scale,<sup>11</sup> which addresses the involuntary muscle activation feature of spasticity<sup>12</sup> as the resistance to a passive movement.<sup>13</sup> The exclusion criteria were as follows: cognitive impairment (score  $>24$  in the Mini-Mental State Examination),<sup>14</sup> diagnosed mood disorder or other neurologic condition, prior severe upper or lower limb trauma, changes in medication for PSS in the previous 48 hours, treatment with botulinum toxin injections within 12 weeks or during the study period, and an epileptic crisis during the previous week. Those in the control group without stroke were recruited from the same population-based cohort.

### Outcome measures

Muscle tone and dynamic stiffness of the biceps brachii and gastrocnemius were assessed with a MyotonPRO.<sup>6,a</sup> The device

contains a probe that applies an initial load of 0.18 N to the skin and then adds up consecutive short impulses (0.40N) to the subcutaneous tissue to characterize mechanical properties. The MyotonPRO calculates muscle tone (tension) by measuring the natural frequency of the acceleration signal and muscle stiffness by measuring the damped natural oscillation response, using an accelerometer.<sup>15</sup> Measurements were taken bilaterally at MB and MT sites, with the muscle relaxed or in the maximum bearable stretched position. The mean score of the 2 consecutive measures was used for the analysis. Regarding the biceps brachii analysis, participants started in relaxed supine position, with the elbow flexed at 45° and forearm in neutral position. For measuring the gastrocnemius, participants lied in prone with approximately 45° of knee flexion. Three sites of muscle testing were included, namely 1 MT location and 2 MB sites. For MB, the mean value at the 2 sites was used in the analysis.

The level of PSS was measured with the MTS,<sup>16</sup> which addresses the muscle response to a manual stretch elicited as slow as possible (V1) and as fast as possible (V3). At fast stretch, muscle tone reflex increases and it is felt at a so-called "catch angle." V1 denotes the passive joint range of movement, whereas V3 denotes the catch angle used to assess spasticity. V1 and V3 were quantified with an electrogoniometer.<sup>b</sup> At V3, the quality of muscle reaction was scored from 0-5, where 0 represents no resistance during passive motion and 5 represents that the joint cannot be moved.<sup>17</sup> To conduct the MTS at the biceps brachii, participants were supine, and the elbow was initially positioned in maximal flexion and supination. For the lower limb, participants remained prone with knees fully extended and feet outside the table.

All outcomes were collected by the same examiner, who had more than 10 years of experience in neurorehabilitation using clinical measures and was previously trained with the MyotonPRO. The examiner remained unaware of the study aims and the participants' allocation group.

### Statistical analysis

Sample size was calculated with the G\*Power software v. 3.1.9.2.<sup>c</sup> We assumed an  $\alpha$  level of 0.05, an 80% statistical power, and a high effect size ( $\eta^2 = 0.15$ ) for differences between groups on muscle tone and stiffness. This generated a sample of 19 participants per group.


Statistical processing was conducted with the PASW Advanced Statistics version 26.0.<sup>d</sup> Normal distribution of the data were evaluated with the Shapiro-Wilk test. We used a mixed-model analysis of variance to compare differences in tone and stiffness of the biceps brachii and gastrocnemius, using muscle position (relaxed vs stretched), side (affected vs nonaffected), and site (MB vs MT) as the within-participant factors and group (subacute stroke, chronic stroke, controls) as the between-participant factor. The Spearman rank test or the Pearson product-moment correlation coefficient analysis were used to test for associations between myotonometry measurement and the MTS. The level of significance was set to  $P < .05$ .

### Data availability

The data that support the study findings are available from the corresponding author on request.

#### List of abbreviations:

<b>MB</b>	<b>muscle belly</b>
<b>MT</b>	<b>musculotendinous</b>
<b>MTS</b>	<b>Modified Tardieu Scale</b>
<b>PSS</b>	<b>poststroke spasticity</b>



**Fig 1** Flowchart diagram of the study participants.

## Results

Seventy-one participants (20 subacute stroke, 20 chronic stroke, 31 controls) were recruited (fig 1). The clinical and demographic characteristics of the sample are listed in table 1.

### Comparison between measurement protocols

Tables 2 and 3 include the tone and stiffness values at the different sites, sides, and groups during the evaluation protocols. The analysis of variance revealed a significant effect of muscle position during myotonometry assessment (relaxed vs stretched) on muscle mechanical properties for (1) the biceps brachii: tone,  $F=59.567$ ,  $P<.001$ ,  $\eta^2=0.095$ ; stiffness,  $F=22.808$ ,  $P<.001$ ,  $\eta^2=0.039$  and (2) the gastrocnemius: tone,  $F=313.2$ ,  $P<.001$ ;  $\eta^2=0.365$ ; stiffness:  $F=341.57$ ;  $P<.001$ ;  $\eta^2=0.386$ . Overall, scores were significantly higher bilaterally and in most testing sites with the muscle stretched than with the muscle relaxed (with a large effect size).

### Discriminative ability between spastic and nonspastic muscles

Myotonometry measurements in relaxed position could not discriminate between the affected and nonaffected sides or between

patients with stroke and controls (all,  $P>.05$ ), except for the lower limb, where higher values were found in the chronic stroke and control groups compared with those with subacute stroke (all,  $P<.05$ ). For assessments in stretched position, differences between sides were only reported at the biceps brachii (increased stiffness in the spastic side,  $P=.020$ ). Furthermore, the comparison between groups demonstrated (1) higher biceps brachii stiffness in the chronic stroke than in the control group ( $P=.045$ ) and (2) lower gastrocnemius tone and stiffness in participants with subacute stroke compared with controls without stroke (all,  $P<.05$ ).

### Discriminative ability between sites of testing

There was a significant muscle position\*sites interaction, with a moderate to large effect size, for (1) the biceps brachii: tone,  $F=8.158$ ,  $P<.004$ ,  $\eta^2=0.015$ ; stiffness,  $F=6.330$ ,  $P<.012$ ,  $\eta^2=0.012$  and (2) the gastrocnemius: tone,  $F=6.089$ ,  $P<.014$ ,  $\eta^2=0.011$ ; stiffness:  $F=39.847$ ,  $P<.001$ ,  $\eta^2=.068$ . Differences between sites of testing were found in the 2 protocols, with higher tone and stiffness at MT sites than MB sites (all  $P<.05$ ), except for the biceps brachii when measured in relaxed position that showed the opposite trend.

**Table 1** Baseline clinical and demographic features of participants

Characteristic	Subacute Stroke (n=20)	Chronic Stroke (n=20)	Control Group (n=31)	P Value
Age (y), mean $\pm$ SD	60.2 $\pm$ 9.7	61.45 $\pm$ 9.7	60.8 $\pm$ 10.6	.926
Sex (female), n, (%)	7 (35)	7 (35)	14 (45.2)	.689
Time after stroke (wk), median (IQR)	17 (6-34)	242.5 (58-1108)	NA	<.001
Affected side, left, n (%)	11 (55)	15 (75)	NA	.289
Hand dominance (right; left; ambidextrous), n (%)	20 (100)	17 (85); 1 (5); 2 (10)	25 (80.6); 6 (19.4)	.131
Leg dominance (right; left; ambidextrous) n (%)	19 (95); 1 (5)	16 (80); 1 (5); 3 (15)	26 (83.9); 5 (16.1)	.319

Abbreviation: NA, Not applicable.

**Table 2** Muscle tone (Hz) and stiffness (N/m) for the biceps brachii at the different sites, sides, and groups during the 2 measurement protocols

Measurement Side		Subacute Stroke Group			Chronic Stroke Group		Control Group	
		Muscle Position	Tone	Stiffness	Tone	Stiffness	Tone	Stiffness
MB sites	Dominant/nonaffected	Relaxed	14.2±1.8	259.1±53.3	14.8±1.5	271.6±40.7	14.7±2.1	271.2±48.7
		Stretched	15.0±1.9*	267.6±50.7	15.3±1.9*	276.4±44.8*	15.7±2.4*	289.3±53.4*
	Nondominant/affected	Relaxed	15.8±2.5	312.3±74.2	15.6±2.1	301.5±60.2	14.6±1.8	276.3±44.6
		Stretched	16.2±2.3	309.2±64.3	16.7±2.7*	315.9±68.3	15.6±2.0*	291.9±43.4*
MT sites	Dominant/nonaffected	Relaxed	14.3±1.3	245.6±33.2	15.1±2.9	275.7±75.2	14.3±1.5	253.4±39.7
		Stretched	16.4±1.8*	285.3±48.7*	16.4±2.3*	290.7±51.8	16.2±1.7*	282.41±36.0*
	Nondominant/affected	Relaxed	14.4±2.5	251.9±54.2	14.2±1.7	259.9±39.1	14.5±1.9	258.4±48.4
		Stretched	16.2±2.3*	285.3±62.7*	16.3±2.5*	307.9±58.7*	16.3±2.1*	288.4±50.1*

\* Significant differences in the within-groups analysis when comparing scores at the same site and side between the 2 different protocols (muscle relaxed vs stretched).

**Table 3** Muscle tone (Hz) and stiffness (N/m) for the gastrocnemius at the different sites, sides, and groups during the 2 measurement protocols

Measurement Side		Subacute Stroke Group			Chronic Stroke Group		Control Group	
		Muscle Position	Tone	Stiffness	Tone	Stiffness	Tone	Stiffness
MB sites	Dominant/nonaffected	Relaxed	14.8±1.3	281.8±28.7	16.4±1.9	291.8±35.8	15.3±1.1	285.6±18.1
		Stretched	18.8±2.6*	348.4±50.8*	20.1±3.4*	387.9±93.7*	19.8±2.2*	383.3±65.2*
	Nondominant/affected	Relaxed	15.1±1.9	283.3±24.7	16.7±2.9	321.0±58.2	15.6±1.4	286.2±21.4
		Stretched	19.0±3.1*	350.2±56.4*	19.7±3.3*	386.6±106.2*	19.8±2.5*	375.4±64.4*
MT sites	Dominant/nonaffected	Relaxed	21.9±2.4	442.2±56.3	22.8±4.0	468.5±94.5	23.9±3.3	483.2±58.6
		Stretched	26.8±4.4*	594.6±123.7*	27.6±3.1*	620.1±103.4*	30.3±4.6*	697.6±128.5*
	Nondominant/affected	Relaxed	21.8±2.9	433.3±63.9	23.0±3.1	459.4±75.9	23.5±2.6	482.5±56.5
		Stretched	27.0±4.7*	577.6±132.8*	27.1±3.3*	599.2±104.2*	28.9±3.7*	659.8±113.5*

\* Significant differences in the within-groups analysis when comparing scores at the same site and side between the 2 different protocols (muscle relaxed vs stretched).

**Table 4** Descriptive data for the clinical measure of spasticity with the Modified Tardieu Scale in the subacute and chronic stroke groups

Measure	Subacute Stroke Group				Chronic Stroke Group			
	V1	V3	V1-V3	X	V1	V3	V1-V3	X
Biceps brachii	172.7±3.6	118.1±6.2	54.5±5.9	2.1±0.1	172.1±3.4	120.0±5.8	52.1±5.3	1.9±0.1
Gastrocnemius	83.1±2.7	65.5±2.4	17.6±2.6	2.3±0.1	78.5±3.5	60.1±4.1	18.4±2.3	2.4±0.1

Abbreviations: V1, joint angle at slow passive stretch (degrees); V3, "catch angle" at fast passive stretch (degrees); X, quality of muscle reaction at V3, from 0-5.

## Correlations

Table 4 lists the clinical data for the measure of spasticity with the MTS in the stroke groups. No significant correlations were observed between myotonometry and the level of PSS, as assessed with the MTS (all  $P > .05$ ).

## Discussion

The present findings partly agree with our hypotheses. Tone and stiffness values changed among the 2 protocols, and myotonometry measurements with the muscle stretched could discriminate between the spastic and nonspastic sides, although only for the biceps brachii. When comparing groups, our results differed depending on the protocol and the assessed muscle. This distinct

behavior has been explained on the basis of the different activation patterns of flexor and extensor muscles.<sup>18</sup>

## Comparison between measurement protocols

Myotonometry is a valid and easy-to-use approach to objectively quantify muscle mechanical properties in people after stroke.<sup>4,6</sup> However, its high environmental sensitivity<sup>4</sup> and the large within- and between-participants variability<sup>19</sup> together with assessment-related aspects, such as muscle position and operator's experience,<sup>4,20</sup> stress the importance of agreeing on a standardized evaluation protocol.

Most previous research in patients with stroke has been conducted carrying out myotonometry measurements with the muscle relaxed. Our findings support the notion that muscle position, relaxed or not, can affect myotonometry scores, which depend on the tissue displacement-force relation.<sup>21</sup> In our study, we mostly

observed higher tone and stiffness during evaluation with the muscle stretched. There are plausible reasons to understand this observation. Motor neuron responsiveness to passive stretch is increased after stroke,<sup>22</sup> which may become more evident with the muscle stretched than relaxed.<sup>23</sup> PSS is also related with shorter muscle fascicles and more compliant tendons that do not respond properly to stretch, increasing tone and stiffness.<sup>24</sup> Additionally, thixotropy, as the influence of movement and time of recovery after movement on mechanical properties,<sup>25</sup> is altered after stroke<sup>26</sup> and can modify muscle stiffness<sup>25</sup> and contribute to intrinsic hypertonia.<sup>27</sup> All in all, changes in mechanical properties after stroke are linked to changes in muscle morphology and composition.<sup>28</sup> This needs to be considered when assessing PSS with myotonometry. It could also explain the lack of association between myotonometry and clinical measures of spasticity, in line with former research<sup>21</sup> but in contradiction with studies that used myotonometry with the muscle contracted.<sup>29,30</sup> The scarce and contradictory literature on this issue, as well as the differences among studies in myotonometry devices and muscle position, can account for the lack of agreement.

### Discriminative ability between spastic and nonspastic muscles

In patients with stroke, myotonometry could only discriminate between sides with the muscle stretched and at the upper limb. In agreement with most literature on the topic,<sup>21,31-33</sup> we observed higher stiffness at the spastic biceps brachii than the nonspastic side. It has been argued that stretching of the biceps brachii evokes higher resistance to elbow extension,<sup>23</sup> and this can make the muscle stiffer and increase tone.<sup>34,35</sup> Nonetheless, evidence on this issue is still preliminary and inconsistent.<sup>36</sup> For the gastrocnemius, myotonometry revealed no differences between sides in any of the protocols. These results agree with previous research using myotonometry to analyze the mechanical properties of different lower limb muscles<sup>33,37-39</sup> in individuals with acute<sup>39</sup> or chronic stroke.<sup>33,37,38</sup> Bilateral adaptations of the lower limbs, especially in those who remain nonphysically active after stroke,<sup>40</sup> could explain the lack of discriminative ability at the gastrocnemius.

Regarding the comparison between spastic and control group muscles, higher tone and stiffness are often expected in chronic poststroke stages,<sup>37,38</sup> although the changes in mechanical properties seem to depend on the assessed muscles.<sup>37,38</sup> As in the present study, biceps brachii tone and stiffness have shown to be increased in patients with chronic stroke.<sup>23,41</sup> Our findings, however, differed for the lower limb, with no differences between the control and chronic stroke groups and with lower tone and stiffness in those with subacute stroke. The reduced stiffness at early stages after stroke has been attributed to a low level of functional recovery.<sup>42</sup> Therefore, the clinical implications may be different for the upper and lower limbs and in patients with different levels of functionality. Future research should include subgroups of participants with different PSS severity and presentation to answer this question.<sup>33</sup>

### Discriminative ability between sites of testing

Current literature suggests that spatial distribution of mechanical properties may not be homogeneous in spastic muscles. Consistent with this, tone and stiffness were significantly different at MT than at MB sites for both muscles and assessment protocols. The

general trend was toward higher tone and stiffness at the tendon, as already observed for the biceps brachii in people with Parkinson disease<sup>43</sup> and for the gastrocnemius in patients with spinal cord injury<sup>44,45</sup> and in healthy volunteers,<sup>46,47</sup> with conflicting evidence for the lower limb.<sup>48</sup> Structural adaptations associated with PSS, for example, lower MB tension with respect to the tendon<sup>49</sup> and lack of muscle strain during stretch<sup>50</sup> and with limb disuse after stroke<sup>51</sup> can help to support these results. Additionally, soft tissue mechanical properties may behave differently, depending on joint position during assessment,<sup>47,52</sup> which highlights again the importance of measuring different spots within the muscle to characterize PSS.<sup>6</sup>

### Study limitations

Several limitations need to be acknowledged. First, the subacute group included patients up to 9 months after stroke, in accordance with previous research on the topic.<sup>9</sup> Despite new standards describing chronic as more than 6 months,<sup>53</sup> endogenous plasticity persists beyond this period,<sup>53</sup> and the chronic stage starts when spontaneous recovery is reduced.<sup>9,54</sup> Therefore, one of the main recommendations for stroke research is to report the time from stroke onset.<sup>53</sup> Second, there was a wide time range after stroke for participants in the chronic group. Third, it could be argued that the MTS would have been more accurate than the Modified Ashworth Scale to screen participants for eligibility.<sup>55</sup> However, in the absence of sufficient psychometric evidence to recommend 1 specific clinical measure,<sup>3</sup> the Modified Ashworth Scale is easy and quick to complete,<sup>13</sup> is highly responsive,<sup>56</sup> and remains the common tool to quantify spasticity after stroke, despite its limitations.<sup>3</sup> Lastly, manual stretch was conducted slowly during evaluation in stretched position to avoid the stretch reflex, although the procedure was not time controlled, and possible muscle activation was not monitored by electromyography. Moreover, the time spent in maximum stretch before evaluation was similar for all participants, but it was not standardized.

### Conclusions

Myotonometry measurements of tone and stiffness can discriminate better between the affected and nonaffected sides in people with stroke and between these and controls without stroke when myotonometry is performed with the muscle stretched. Clinical measures of spasticity were not correlated with myotonometry, regardless of the muscle position during evaluation.

### Suppliers

- a. MyotonPRO; Myoton AS.
- b. Electrogoniometer; Biometrics Ltd.
- c. G\*Power software v. 3.1.9.2; Heinrich-Heine University.
- d. PASW Advanced Statistics version 26.0; SPSS Inc.

### Keywords

Muscle spasticity; Outcome assessment, health care; Rehabilitation; Stroke

## Corresponding author

Paula González-García, PhD, Departamento de Fisioterapia, Facultad de Enfermería, Fisioterapia y Podología, Universidad de Sevilla, c/ Avicena s/n, 41009 Sevilla, Spain. *E-mail address:* pgonzalez@us.es.

## Acknowledgments

We thank all the participants who took part in the study, and we thank the stroke units at CRECER and DACE for helping us during the recruitment process.

## References

- Zeng H, Chen J, Guo Y, Tan S. Prevalence and risk factors for spasticity after stroke: a systematic review and meta-analysis. *Front Neurol* 2021;11:616097.
- Lehoux MC, Sobczak S, Cloutier F, Charest S, Bertrand-Grenier A. Shear wave elastography potential to characterize spastic muscles in stroke survivors: literature review. *Clin Biomech* 2020;72:84–93.
- Aloraini SM, Gäverth J, Yeung E, MacKay-Lyons M. Assessment of spasticity after stroke using clinical measures: a systematic review. *Disabil Rehabil* 2015;37:2313–23.
- Luo Z, Lo WLA, Bian R, Wong S, Li L. Advanced quantitative estimation methods for spasticity: a literature review. *J Int Med Res* 2019;48:300060519888425.
- Tran A, Gao J. Quantitative ultrasound to assess skeletal muscles in post stroke spasticity. *J Cent Nerv Syst Dis* 2021; 13:1179573521996141.
- García-Bernal MI, Heredia-Rizo AM, González-García P, Cortés-Vega MD, Casuso-Holgado MJ. Validity and reliability of myotonometry for assessing muscle viscoelastic properties in patients with stroke: a systematic review and meta-analysis. *Sci Rep* 2021; 11:5062.
- Megna M, Marvulli R, Farì G, et al. Pain and muscles properties modifications after botulinum toxin type A (BTX-A) and radial extracorporeal shock wave (rESWT) combined treatment. *Endocr Metab Immune Disord Drug Targets* 2019;19:1127–33.
- Burridge JH, Wood DE, Hermens HJ, et al. Theoretical and methodological considerations in the measurement of spasticity. *Disabil Rehabil* 2005;27:69–80.
- Chuang LL, Wu CY, Lin KC, Lur SY. Quantitative mechanical properties of the relaxed biceps and triceps brachii muscles in patients with subacute stroke: a reliability study of the Myoton-3 myometer. *Stroke Res Treat* 2012;2012:617694.
- Sarasso S, Määttä S, Ferrarelli F, Poryazova R, Tononi G, Small SL. Plastic changes following imitation-based speech and language therapy for aphasia: a high-density sleep EEG study. *Neurorehabil Neural Repair* 2014;28:129–38.
- Ansari N, Naghdi S, Arab T, Jalair S. The interrater and intrarater reliability of the Modified Ashworth Scale in the assessment of muscle spasticity: limb and muscle group effect. *NeuroRehabilitation* 2008;23:231–7.
- Shu X, McConaghy C, Knight A. Validity and reliability of the Modified Tardieu Scale as a spasticity outcome measure of the upper limbs in adults with neurological conditions: a systematic review and narrative analysis. *BMJ Open* 2021;11:1–10.
- Bohannon R, Smith M. Interrater reliability of a Modified Ashworth Scale of muscle spasticity. *Phys Ther* 1987;67:206–7.
- Quinn TJ, Elliott E, Langhorne P. Cognitive and mood assessment tools for use in stroke. *Stroke* 2018;49:483–90.
- Ilahi S, T Masi A, White A, Devos A, Henderson J, Nair K. Quantified biomechanical properties of lower lumbar myofascia in younger adults with chronic idiopathic low back pain and matched healthy controls. *Clin. Biomech* 2020;73:78–85.
- Paulis WD, Horemans HLD, Brouwer BS, Stam HJ. Excellent test-retest and inter-rater reliability for Tardieu Scale measurements with inertial sensors in elbow flexors of stroke patients. *Gait Posture* 2011;33:185–9.
- Petek Balci B. Spasticity measurement. *Noro Psikiyatr Ars* 2018;55 (Suppl 1):S49–53.
- Puce L, Currà A, Marinelli L, et al. Clinical neurophysiology practice spasticity, spastic dystonia, and static stretch reflex in hypertonic muscles of patients with multiple sclerosis. *Clin Neurophysiol Pract* 2021;6:194–202.
- Eby SF, Zhao H, Song P, et al. Quantifying spasticity in individual muscles using shear wave elastography. *Radiol Case Reports* 2017;12:348–52.
- Amirova LE, Plehuna A, Rukavishnikov IV, Saveko AA, Peipsi A, Tomilovskaya ES. Sharp changes in muscle tone in humans under simulated microgravity. *Front Physiol* 2021;12:1–12.
- Xiaoyan L, Shin H, Zong Y, Li S, Zhou P. Assessing muscle compliance in stroke with the myotonometer. *Clin Biomech* 2017;50:110–3.
- Condliffe EG, Clark DJ, Patten C. Reliability of elbow stretch reflex assessment in chronic post-stroke hemiparesis. *Clin Neurophysiol* 2005;116:1870–8.
- Gao J, He W, Du LJ, et al. Quantitative ultrasound imaging to assess the biceps brachii muscle in chronic post-stroke spasticity: preliminary observation. *Ultrasound Med Biol* 2018;44:1931–40.
- Davis JF, Khir AW, Barber L, et al. The mechanisms of adaptation for muscle fascicle length changes with exercise: implications for spastic muscle. *Med Hypotheses* 2020;144:110199.
- Lakie M, Campbell KS. Muscle thixotropy—where are we now? *J Appl Physiol* 2019;126:1790–9.
- Vattanasilp W, Ada L, Crosbie J. Contribution of thixotropy, spasticity, and contracture to ankle stiffness after stroke. *J Neurol Neurosurg Psychiatry* 2000;69:34–9.
- Bakheit AMO. The pharmacological management of post-stroke muscle spasticity. *Drugs Aging* 2012;29:941–7.
- Lieber RL, Ward SR. Cellular mechanisms of tissue fibrosis. 4. Structural and functional consequences of skeletal muscle fibrosis. *Am J Physiol Cell Physiol* 2013;305:C241–52.
- Leonard CT, Stephens JU, Stroppel SL. Assessing the spastic condition of individuals with upper motoneuron involvement: validity of the myotonometer. *Arch Phys Med Rehabil* 2001;82:1416–20.
- Rydahl SJ, Brouwer BJ. Ankle stiffness and tissue compliance in stroke survivors: a validation of myotonometer measurements. *Arch Phys Med Rehabil* 2004;85:1631–7.
- Li X, Shin H, Li S, Zhou P. Assessing muscle spasticity with myotonometric and passive stretch measurements: validity of the myotonometer. *Sci Rep* 2017;7:1–7.
- Leng Y, Lo WLA, Hu C, et al. The effects of extracorporeal shock wave therapy on spastic muscle of the wrist joint in stroke survivors: evidence from neuromechanical analysis. *Front Neurosci* 2021;14:1–16.
- Miller T, Ying MTC, Chung RCK, Pang MYC. Convergent validity and test-retest reliability of multimodal ultrasonography and related clinical measures in people with chronic stroke. *Arch Phys Med Rehabil* 2022;103:459–72.
- Wu CH, Ho YC, Hsiao MY. Evaluation of post-stroke spastic muscle stiffness using shear wave ultrasound elastography. *Ultrasound Med Biol* 2017;43:1105–11.
- Haugh A, Pandyan A, Johnson G. A systematic review of the Tardieu Scale for the measurement of spasticity. *Disabil Rehabil* 2006;28:899–907.
- Chuang LL, Wu CY, Lin KC. Reliability, validity, and responsiveness of myotonometric measurement of muscle tone, elasticity, and stiffness in patients with stroke. *Arch Phys Med Rehabil* 2012;93:532–40.
- Fröhlich-Zwahlen AK, Casartelli NC, Item-Glatthorn JF, Maffiuletti NA. Validity of resting myotonometric assessment of lower extremity muscles in chronic stroke patients with limited hypertonia: a preliminary study. *J Electromyogr Kinesiol* 2014;24:762–9.

38. Wang JS, Lee SB, Moon SH. The immediate effect of PNF pattern on muscle tone and muscle stiffness in chronic stroke patient. *J Phys Ther Sci* 2016;28:967–70.
39. Lo WLA, Zhao JL, Li L, Mao YR, Huang DF. Relative and absolute interrater reliabilities of a hand-held myotonometer to quantify mechanical muscle properties in patients with acute stroke in an inpatient ward. *Biomed Res Int* 2017;2017:4294028.
40. Berenpas F, Martens AM, Weerdesteyn V, Geurts AC, van Alfen N. Bilateral changes in muscle architecture of physically active people with chronic stroke: a quantitative muscle ultrasound study. *Clin Neurophysiol* 2017;128:115–22.
41. Lee S, Spear S, Rymer W. Quantifying changes in material properties of stroke-impaired muscle. *Clin Biomech* 2015;30:269–75.
42. Mirbagheri MM, Tsao C, Rymer WZ. Natural history of neuromuscular properties after stroke: a longitudinal study. *J Neurol Neurosurg Psychiatry* 2009;80:1212–7.
43. Marusiak J, Jaskólska A, Budrewicz S, Koszewicz M, Jaskólski A. Increased muscle belly and tendon stiffness in patients with Parkinson's disease, as measured by myotonometry. *Mov Disord* 2011;26:2119–22.
44. Ge JS, Chang TT, Zhang ZJ. Reliability of myotonometric measurement of stiffness in patients with spinal cord injury. *Med Sci Monit* 2020;26:1–7.
45. Ko CY, Choi HJ, Ryu J, Kim G. Between-day reliability of MyotonPRO for the non-invasive measurement of muscle material properties in the lower extremities of patients with a chronic spinal cord injury. *J Biomech* 2018;73:60–5.
46. Feng YN, Li YP, Liu CL, Zhang ZJ. Assessing the elastic properties of skeletal muscle and tendon using shearwave ultrasound elastography and MyotonPRO. *Sci Rep* 2018;8:1–9.
47. Huang J, Qin K, Tang C, et al. Assessment of passive stiffness of medial and lateral heads of gastrocnemius muscle, achilles tendon, and plantar fascia at different ankle and knee positions using the myotonPRO. *Med Sci Monit* 2018;24:7570–6.
48. Theis N, Mohagheghi AA, Korff T. Mechanical and material properties of the plantarflexor muscles and Achilles tendon in children with spastic cerebral palsy and typically developing children. *J Biomech* 2016;49:3004–8.
49. Faturi FM, Lopes Santos G, Ocamoto GN, Russo TL. Structural muscular adaptations in upper limb after stroke: a systematic review. *Top Stroke Rehabil* 2019;26:73–9.
50. Chardon MK, Suresh NL, Dhaher YY, Rymer WZ. In-vivo study of passive musculotendon mechanics in chronic hemispheric stroke survivors. *IEEE Trans Neural Syst Rehabil Eng* 2020;28:1022–31.
51. Narici MV, Maganaris CN. Plasticity of the muscle-tendon complex with disuse and aging. *Exerc Sport Sci Rev* 2007;35:126–34.
52. Chang TT, Feng YN, Zhu Y, Liu CL, Wang XQ, Zhang ZJ. Objective assessment of regional stiffness in achilles tendon in different ankle joint positions using the MyotonPRO. *Med Sci Monit* 2020;26:1–8.
53. Bernhardt J, Hayward KS, Kwakkel G, et al. Agreed definitions and a shared vision for new standards in stroke recovery research: the Stroke Recovery and Rehabilitation Roundtable taskforce. *Int J Stroke* 2017;12:444–50.
54. Nichols-Larsen DS, Clark PC, Zeringue A, Greenspan A, Blanton S. Factors influencing stroke survivors' quality of life during subacute recovery. *Stroke* 2005;36:1480–4.
55. Patrick E, Ada L. The Tardieu Scale differentiates contracture from spasticity whereas the Ashworth Scale is confounded by it. *Clin Rehabil* 2006;20:173–81.
56. Chen CL, Chen CY, Chen HC, et al. Responsiveness and minimal clinically important difference of Modified Ashworth Scale in patients with stroke. *Eur J Phys Rehabil Med* 2019;55:754–60.